

VACCINATION GUIDELINES FOR FOALS

NON-VACCINATED DAM	VACCINATED DAM
3 Months: 3- Way, Rabies, WNV	6 Months: 3-Way, Rabies, Rhino, WNV
4 Months: 3-Way, WNV	7 Months: 3-Way, Rabies, Rhino, WNV
5 Months: 3-Way, Rhino, WNV	8-9 Months: 4-Way, Rhino, WNV
6 Months: Influenza/Rhino	10 Months: Influenza/Rhino
7 Months: Influenza/Rhino	11-12 Months: Rabies, Influenza I/N
8 Months: Influenza I/N	Strangles Injection: 6th, 7th, 8th and 12th month, then every 6 months (may want to use I/N 11th or 12th month).
12 Months: Rabies	Strangles I/N: 11th & 12th month, then every 6 months (start at 6th months for endemic areas).
Strangles Injection: 6th, 7th, 8th and 12th month, then every 6 months (may want to use I/N 11th or 12th month).	3-Way: (E/W Encephalomyelitis, Tetanus Toxoid).
Strangles I/N: 11th & 12th month, then every 6 months (start at 6th months for endemic areas).	4-Way: (E/W Encephalomyelitis, Influenza, Tetanus Toxoid).
3-Way: (E/W Encephalomyelitis, Tetanus Toxoid).	Rhinopneumonitis: (EHV-4 The form of this virus that causes respiratory disease).
Rhinopneumonitis: (EHV-4 The form of this virus that causes respiratory disease).	Equine Herpes Virus:
Equine Herpes Virus:	EHV-1: The form of this virus that primarily causes abortion. Start at 4 months if in high-risk area.
EHV-1: The form of this virus that primarily causes abortion.	EHV-4: Certain strains can cause neurological disease.
EHV-4: Certain strains can cause neurological disease.	De-Worming: Start at 2 months and de-worm every 30 days until 8 months of age. Then continue to de-worm every 60 days. Use Ivermectin, Strongid or Panacur.
De/Worming: Start at 2 months and de-worm every 30 days until 8 months of age. Then continue to de-worm every 60 days. Use Ivermectin, Strongid or Panacur.	

THIS IS A SUGGESTED VACCINATION SCHEDULE ONLY. THE VETERINARIAN MAY CHANGE THIS SCHEDULE BASED ON INDIVIDUAL NEEDS, FARM/RANCH MANAGEMENT, OR CURRENT DISEASE OUTBREAK.